



RSC96-CopGFP Cell Line User Guide

► Basic Product Information

GFP (Green Fluorescent Protein) is a vital tool for observing molecular dynamics within cells. However, it often encounters challenges such as insufficient brightness and limited stability in expression patterns. To tackle these issues, EDITGENE has developed the OE-Booster cis-regulatory element, which enables the production of high-brightness and high-stability CopGFP-tagged cell lines. These cell lines are versatile for use in multiplex fluorescence labeling experiments, enhancing the accuracy of intracellular dynamic observations and improving molecular tracking to reveal complex biological processes.

The CopGFP stable cell lines from EDITGENE are constructed using lentiviral transfection, ensuring stable and efficient expression of the CopGFP fluorescent protein. These cell lines can serve as control lines in lentiviral infection experiments.

Product Name	RSC96-CopGFP
Catalog No.	EDC070-CopGFP
Cell Morphology	Adherent
Trypsinization Time	30 s~1 min
Fluorescence	-
Resistance	Puromycin
Drug Administration	Puro = 1.5 µg/mL
Passage Ratio	1:10
Complete Culture Medium	DMEM(GIBCO:C11995500BT)+10%FBS(GIBCO:FBS-S500)
Cryopreservation Culture Medium	50% basic culture medium+40%FBS+10%DMSO (SIGMA : D2650-100ML)



► Advantages of CopGFP Cell Line:

Exceptional Brightness

EDITGENE CopGFP cell lines are optimized to significantly enhance fluorescence intensity, enabling clearer observation of dynamic cellular processes and capturing even the smallest changes.

Outstanding Stability

Specifically designed to ensure consistency over long-term experiments, these cell lines minimize fluorescence decay, providing more reliable experimental results.


Broad Applicability

Suitable for various cell types, including cancer cells and stem cells, these cell lines meet diverse research needs under different experimental conditions.

Efficient Expression

Associated with OE-Booster technology, EDITGENE CopGFP cell lines achieve a remarkable improvement in protein expression efficiency, enhancing the effectiveness of your experiments.


► Plasmid Map






Verification Images

bright field image



fluorescent image



Cell Receipt

a. Frozen Cells

If the cells are shipped on dry ice, immediately transfer them to liquid nitrogen storage or proceed with cell thawing directly upon receipt.

b. Live Cells

Upon receipt, disinfect the external surface of the T25 flask with 75% ethanol. Place the flask in a 5% CO₂, 37°C incubator for 2 hours. After incubation, observe the cell attachment and confluency under a microscope. Take two photos at different fields of view at 100x and 40x magnification. If the confluency is greater than 80%, proceed with passaging. If the confluency is less than 80%, discard the medium in the flask, replace it with fresh complete medium, and continue incubation.

Note: Do not overfill the culture flask with medium, as it may affect cell culture conditions.

Cell Thawing Procedure

1. Preheat the water bath to 37°C.
2. Prewarm the complete medium suitable for this cell line to 37°C.
3. Prepare a 15 mL centrifuge tube with 6 mL of complete medium.
4. Retrieve the frozen cells from liquid nitrogen and gently swirl the cryovial in a 37°C water bath until only a small ice core remains. Thaw the cells within 2 minutes.

Note: Do not submerge the vial cap in water or seal the cap with parafilm.

5. Transfer the cryovial to a sterile environment. Disinfect the outside of the cryovial with 75% ethanol before opening.
6. Using a pipette, transfer the cell suspension into the prewarmed complete medium in the centrifuge





tube.

7. Centrifuge the cell suspension at 500 g for 4 minutes.
8. After centrifugation, check the supernatant for clarity and verify the presence of a pellet. Carefully aspirate the supernatant under sterile conditions, and gently resuspend the pellet in 1 mL of complete medium. Ensure thorough mixing by gentle pipetting.
9. Seed the cells into a T25 culture flask or an equivalent surface area container, adding 4 mL of complete medium.
10. Gently swirl the flask to ensure even cell distribution and incubate at 37°C, 5% CO₂ (the specific environment depends on the cell type and culture medium used).
11. Observe cell status the next day.
 - For adherent cells: If cells are well attached, replace with fresh medium. If cells appear rounded but unattached, allow another 24 hours before changing the medium. Subsequently, change the medium every 2-3 days based on cell growth and passage when 80% confluency is reached.
 - For suspension cells: If cell viability is good, replace with fresh medium. If cells appear unhealthy or gray, observe for an additional 24 hours before proceeding with medium changes.

► Cell Passaging

a. Adherent Cells

1. Prewarm complete medium, PBS, and trypsin to 37°C.
2. Aspirate the supernatant from the culture vessel.
3. Gently add PBS to one side of the vessel (approximately 2 mL for a T25 flask) to wash the cells. Ensure gentle washing and thorough coverage without disturbing the cell layer. Aspirate the PBS by tilting the flask back and forth.
4. Add 1 mL of trypsin to the T25 flask and distribute evenly. Place the flask in an incubator for digestion.
5. Observe under a microscope for cell detachment (70%-80% cells should round up). Gently tap the flask to detach cells from the surface.
6. Immediately add 2-3 times the volume of complete medium to neutralize the trypsin and gently mix to stop digestion.
7. Collect the cell suspension and pipette the bottom of the flask to ensure all cells are detached. Avoid creating bubbles that could damage cells.
8. Centrifuge the collected cell suspension at 500 g for 5 minutes.





9. After centrifugation, aspirate the supernatant, resuspend the pellet in 1 mL of complete medium, and gently mix.
10. Plate the cells at the appropriate ratio. For the first passage, we recommend a 1:2 ratio. If the cells reach confluency within two days, you can increase the ratio. If it takes 3-4 days to reach confluency, reduce the passaging ratio accordingly.
11. Gently swirl the flask and incubate at 37°C, 5% CO₂ (for vented flasks, loosen the cap before placing in the incubator to allow proper gas exchange).
12. Observe cell status the next day. If a large number of dead cells are present, change the medium. Afterward, change the medium daily based on cell growth until they reach 80% confluency, at which point passaging or cryopreservation is necessary.

b. Suspension Cells

1. Prewarm complete medium, PBS, and trypsin to 37°C.
2. Collect the cell suspension and pipette the bottom of the flask to ensure all cells are detached. Avoid creating bubbles that could damage cells.
3. Centrifuge the collected cell suspension at 500 g for 4 minutes.
4. After centrifugation, aspirate the supernatant, resuspend the pellet in 1 mL of complete medium, and gently mix.
5. Plate the cells at the appropriate ratio. For the first passage, we recommend a 1:2 ratio. If the cells reach confluency within two days, you can increase the ratio. If it takes 3-4 days to reach confluency, reduce the passaging ratio accordingly.
6. Gently swirl the flask and incubate at 37°C, 5% CO₂ (for vented flasks, loosen the cap before placing in the incubator to allow proper gas exchange).
7. Observe cell status the next day. If a large number of dead cells are present, change the medium. Afterward, change the medium daily based on cell growth until they reach 80% confluency, at which point passaging or cryopreservation is necessary.

c. Semi-adherent and Semi-suspension Cells

1. Prewarm complete medium, PBS, and trypsin to 37°C.
2. Collect the cell suspension and pipette the bottom of the flask to ensure all cells are detached. Avoid creating bubbles that could damage cells.
3. Gently add PBS to one side of the vessel (approximately 2 mL for a T25 flask) to wash the cells.





Ensure gentle washing and thorough coverage without disturbing the cell layer. Aspirate the PBS by tilting the flask back and forth.

4. Add 1 mL of trypsin to the T25 flask and distribute evenly. Place the flask in an incubator for digestion.
5. Observe under a microscope for cell detachment (70%-80% cells should round up). Gently tap the flask to detach cells from the surface.
6. Immediately add 2-3 times the volume of complete medium to neutralize the trypsin and gently mix to stop digestion.
7. Collect the cell suspension and pipette the bottom of the flask to ensure all cells are detached. Avoid creating bubbles that could damage cells.
8. Centrifuge the collected cell suspension at 500 g for 5 minutes.
9. After centrifugation, aspirate the supernatant, resuspend the pellet in 1 mL of complete medium, and gently mix.
10. Plate the cells at the appropriate ratio. For the first passage, we recommend a 1:2 ratio. If the cells reach confluency within two days, you can increase the ratio. If it takes 3-4 days to reach confluency, reduce the passaging ratio accordingly.
11. Gently swirl the flask and incubate at 37°C, 5% CO₂ (for vented flasks, loosen the cap before placing in the incubator to allow proper gas exchange).
12. Observe cell status the next day. If a large number of dead cells are present, change the medium. Afterward, change the medium daily based on cell growth until they reach 80% confluency, at which point passaging or cryopreservation is necessary.

Note: To maintain the stable expression of Luciferase gene, culture with drug administration is recommended.

► Cell Cryopreservation

1. Collect cells as per the passaging protocol and adjust the volume of medium based on the size of the pellet.
2. Gently resuspend the pellet in cold cryopreservation medium.
3. Take 20 µL of the suspension for cell counting.
4. Centrifuge at 500 g for 5 minutes at room temperature.
5. Aspirate the supernatant, then resuspend the pellet in 1-2 mL of pre-cooled cryopreservation medium to achieve a density of 1×10^6 cells/mL.





6. Aliquot the cell suspension into cryovials (1 mL per vial) and label each vial with the cell name, passage number, cell count, and cryopreservation date.
 7. Place the cryovials in a pre-cooled freezing container at 4°C and transfer them to an ultra-low temperature freezer within 15 minutes.
 8. After overnight freezing, transfer the cryovials to liquid nitrogen storage for long-term preservation.
-

Precautions

- Upon receiving room temperature cells, inspect the vial for any leakage or damage, and document accordingly.
- Disinfect the culture flask with 75% ethanol and observe the cell condition under a microscope. Place the flask in a cell culture incubator to stabilize for 2-4 hours before handling.
- Carefully review the user manual to familiarize yourself with the cell line, including adherence properties, morphology, base medium, passaging ratios, and frequency of medium changes.
- After stabilization, remove the flask and inspect the cells again. Photograph and record the cell condition, as this will be used for follow-up services. We recommend photographing the cells during each passage to document growth status.
- If you notice any abnormalities or have questions about the cells, please contact our support team for assistance.

Advantages



Species Diversity

Over 100 types of wild-type cells from species including human, mouse, chicken, pig, and cow, covering a wide range of research areas.



STR Authentication

Each cell line undergoes STR/species authentication and stringent quality control to ensure cell identity.



Experimental Validation



All cell lines in this collection have been validated for gene-editing experiments, making them suitable for most gene-editing applications.



Authoritative Source

All cell lines are sourced from reputable cell banks such as ATCC and the Chinese Academy of Sciences, ensuring low passage, high viability, and optimal cell status.

